

On the upper Assouad spectrum

ALEX RUTAR

ABSTRACT. The goal of this note is to provide a simplified proof of the following fact: if E is a non-empty metric space with $\dim_{\text{qA}} E < \infty$, then for $0 \leq \theta < 1$,

$$\overline{\dim}_A^\theta E = \sup_{0 \leq \lambda \leq \theta} \dim_A^\lambda E.$$

This will follow from an asymptotic Lipschitz property of a two-scale branching function associated with the set E .

This result was previously obtained in [FHH+19] under the additional assumption that E is a subset of Euclidean space, with a more difficult proof.

A two-scale branching function. Let E be a non-empty metric space. We first recall the *two-scale branching function* associated with E , which was introduced in [OR25+]. It is the function β defined for $0 \leq v \leq u$ by

$$\beta(u, v) = \log \sup_{x \in E} N_{2^{-u}}(B(x, 2^{-v})).$$

Here, the base of the logarithm is 2; for $\delta > 0$, $B(x, \delta)$ denotes the open ball with radius δ ; and $N_\delta(F)$ denotes the least number of open balls of radius δ required to cover a set $F \subset E$.

We begin with some basic properties of the function β .

Lemma 1. *Let E have two-scale branching function β . Then:*

- (i) $\beta(u, u) = 0$ for all $u \geq 0$.
- (ii) $\beta(u, v)$ is increasing in u and decreasing in v .
- (iii) For all $0 \leq v \leq w \leq u$,

$$(1) \quad \beta(u, v) \leq \beta(u, w) + \beta(w, v).$$

Proof. The fact that $\beta(u, u) = 0$ for $u \geq 0$ is immediate, along with the monotonicity properties. To verify (1), let $0 \leq v \leq w \leq u$ be arbitrary. Then to obtain a cover for $B(x, 2^{-v})$, we first cover with balls of radius 2^{-w} , and then cover each ball in the resulting cover with balls of radius 2^{-u} . Therefore

$$N_{2^{-u}}(B(x, 2^{-v})) \leq N_{2^{-w}}(B(x, 2^{-v})) \cdot \sup_{y \in E} N_{2^{-u}}(B(y, 2^{-w})).$$

Taking suprema in x and a logarithm, the claim follows. □

Assouad spectra and quasi-Assouad dimension. Using the two-scale branching function β , let us recall various definitions associated with the Assouad spectrum. First, the *Assouad spectrum* is defined for $0 \leq \theta < 1$ by

$$\dim_A^\theta E = \inf \left\{ s \geq 0 : \exists C \geq 0 \forall 0 \leq u \text{ we have } \beta(u, \theta u) \leq C + s(1 - \theta)u \right\}.$$

Similarly, the *upper Assouad spectrum* is defined for $0 \leq \theta < 1$ by

$$\overline{\dim}_A^\theta E = \inf \left\{ s \geq 0 : \exists C \geq 0 \forall 0 \leq v \leq \theta u \text{ we have } \beta(u, v) \leq C + s(u - v) \right\}.$$

Clearly $\overline{\dim}_A^\theta E \geq \dim_A^\theta E$, and moreover $\overline{\dim}_A^\theta E$ is monotonically increasing in θ . The *quasi-Assouad dimension* is the limit at 1:

$$\dim_{\text{qA}} E = \lim_{\theta \nearrow 1} \overline{\dim}_A^\theta E.$$

It is standard to rephrase the above definitions in terms of limits of β .

Lemma 2. *Let E have two-scale branching function β . Then*

$$\dim_A^\theta E = \limsup_{u \rightarrow \infty} \frac{\beta(u, \theta u)}{u(1 - \theta)}$$

and

$$\overline{\dim}_A^\theta E = \limsup_{u \rightarrow \infty} \sup_{0 \leq \lambda \leq \theta} \frac{\beta(u, \lambda u)}{u(1 - \lambda)}.$$

We can now see that the result claimed in the abstract is simply a justification of the exchange of the limit and the supremum in the equivalent formula for $\overline{\dim}_A^\theta E$. To do this, we will prove that the functions $\beta(u, v)$ are uniformly Lipschitz in the variable v up to a sub-linear error term in u .

An asymptotic Lipschitz property. Now let us assume in addition that the quasi-Assouad dimension of E is finite.

Lemma 3. *Let E have two-scale branching function β , and suppose $\alpha = \dim_{\text{qA}} E < \infty$. Then for all $0 \leq v \leq u$,*

$$\beta(u, v) \leq \alpha(u - v) + o(u).$$

Proof. It suffices to show for all $\varepsilon > 0$ there is a constant $C_\varepsilon \geq 1$ such that for all $0 < r \leq R < 1$,

$$(2) \quad \sup_{x \in E} N_r(B(x, R)) \leq C_\varepsilon r^{-\varepsilon} \left(\frac{R}{r} \right)^\alpha.$$

Let $\theta < 1$ be sufficiently large so that $(1 - \theta)(\alpha + \varepsilon) \leq \varepsilon$. We consider two cases depending on the value of R . If $R \geq r^\theta$, then by definition of the quasi-Assouad

dimension there is a constant C_ε (depending on E , θ , and ε) so that

$$N_r(B(x, R)) \leq C_\varepsilon \left(\frac{R}{r}\right)^{\alpha+\varepsilon} \leq C_\varepsilon r^{-\varepsilon} \left(\frac{R}{r}\right)^\alpha.$$

In the second inequality we just use that $R < 1$. Otherwise, if $R \leq r^\theta$, since $B(x, R) \subset B(x, r^\theta)$,

$$N_r(B(x, R)) \leq N_r(B(x, r^\theta)) \leq C_\varepsilon \left(\frac{r^\theta}{r}\right)^{\alpha+\varepsilon} \leq C_\varepsilon r^{-(1-\theta)(\alpha+\varepsilon)} \leq C_\varepsilon r^{-\varepsilon} \left(\frac{R}{r}\right)^\alpha$$

where the last line follows since $(R/r)^\alpha \geq 1$. Since $x \in E$ was arbitrary, the claim in (2) follows.

Taking logarithms and substituting the definition of β , the desired claim follows. $\square$

Using the subadditivity property (1), we can convert Lemma 3 into an asymptotic Lipschitz property in the second argument of β .

Corollary 4. *Let E have two-scale branching function β , and suppose $\alpha = \dim_{\text{qA}} E < \infty$. Then for all $0 \leq w \leq v \leq u$,*

$$0 \leq \beta(u, w) - \beta(u, v) \leq \alpha(v - w) + o(u).$$

Proof. By monotonicity, $\beta(u, w) \geq \beta(u, v)$. Then by (1) followed by Lemma 3,

$$\beta(u, w) - \beta(u, v) \leq \beta(w, v) \leq \alpha(v - w) + o(u)$$

as claimed. $\square$

Upper Assouad spectra from Assouad spectra. Now, we prove that we can recover the upper Assouad spectrum from the Assouad spectrum.

Theorem 5. *Let E be a non-empty metric space with $\dim_{\text{qA}} E < \infty$. Then for $0 \leq \theta < 1$,*

$$\overline{\dim}_A^\theta E = \sup_{0 \leq \lambda \leq \theta} \dim_A^\lambda E.$$

Proof. Recalling Lemma 2, get an increasing sequence $0 < u_n \rightarrow \infty$ and $\lambda_n \in [0, \theta]$ such that $\lambda_n \rightarrow \lambda \in [0, \theta]$ and

$$\overline{\dim}_A^\theta E = \lim_{n \rightarrow \infty} \frac{\beta(u_n, \lambda_n u_n)}{u_n(1 - \lambda_n)}.$$

Then by Corollary 4, for $n \in \mathbb{N}$,

$$(3) \quad |\beta(u_n, \lambda_n u_n) - \beta(u_n, \lambda)| \leq \alpha u_n |\lambda_n - \lambda| + o(u_n).$$

Since $\theta < 1$, $(1 - \lambda_n)^{-1} \rightarrow (1 - \lambda)^{-1}$, so

$$\lim_{n \rightarrow \infty} \left| \frac{\beta(u_n, \lambda_n u_n)}{u_n(1 - \lambda_n)} - \frac{\beta(u_n, \lambda)}{u_n(1 - \lambda)} \right| = 0.$$

Therefore $\dim_A^\lambda E \geq \overline{\dim}_A^\theta E$ as required. $\square$

ACKNOWLEDGEMENTS

The author thanks Vilma Orgoványi for interesting discussions around the contents of this note.

REFERENCES

- [FHH+19] J. M. Fraser, K. E. Hare, K. G. Hare, S. Troscheit, and H. Yu. *The Assouad spectrum and the quasi-Assouad dimension: a tale of two spectra*. Ann. Acad. Sci. Fenn., Math. **44** (2019), 379–387. [zb1:1410.28008](#).
- [OR25+] V. Orgoványi and A. Rutar. *Two-scale branching functions and inhomogeneous attractors*. Preprint. [arxiv:2510.07013v1](#).

Alex Rutar

Department of Mathematics and Statistics, University of Jyväskylä, P.O. Box 35 (MaD), FI-40014 University of Jyväskylä, Finland

Email: alex@rutar.org